

X757/75/02

Physics Section 1—Questions

TUESDAY, 5 MAY 9:00 AM - 11:00 AM

Instructions for the completion of Section 1 are given on *Page two* of your question and answer booklet X757/75/01.

Record your answers on the answer grid on *Page three* of your question and answer booklet.

Reference may be made to the Data Sheet on *Page two* of this booklet and to the Relationship Sheet X757/75/11.

Before leaving the examination room you must give your question and answer booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





Speed of light in materials

Material	Speed in m s ⁻¹
Air	$3.0 imes 10^8$
Carbon dioxide	$3.0 imes 10^8$
Diamond	1.2×10^8
Glass	2.0×10^8
Glycerol	$2 \cdot 1 \times 10^8$
Water	$2\cdot3 imes10^8$

Gravitational field strengths

	Gravitational field strength on the surface in N kg ⁻¹
Earth	9.8
Jupiter	23
Mars	3.7
Mercury	3.7
Moon	1.6
Neptune	11
Saturn	9.0
Sun	270
Uranus	8.7
Venus	8.9

Specific latent heat of fusion of materials

Material	Specific latent heat of fusion in Jkg ⁻¹
Alcohol	0.99×10^5
Aluminium	$3.95 imes 10^5$
Carbon Dioxide	1.80×10^5
Copper	2.05×10^5
Iron	$2 \cdot 67 \times 10^5$
Lead	0.25×10^5
Water	3.34×10^5

Specific latent heat of vaporisation of materials

Material	Specific latent heat of vaporisation in J kg ⁻¹
Alcohol	11.2×10^5
Carbon Dioxide	3.77×10^5
Glycerol	$8\cdot 30 imes 10^5$
Turpentine	$2.90 imes 10^5$
Water	22.6 $\times 10^5$

Speed of sound in materials

	1
Material	Speed in m s ⁻¹
Aluminium	5200
Air	340
Bone	4100
Carbon dioxide	270
Glycerol	1900
Muscle	1600
Steel	5200
Tissue	1500
Water	1500

Specific heat capacity of materials

Material	Specific heat capacity in J kg ⁻¹ °C ⁻¹			
Alcohol	2350			
Aluminium	902			
Copper	386			
Glass	500			
Ice	2100			
Iron	480			
Lead	128			
Oil	2130			
Water	4180			

Melting and boiling points of materials


Material	Melting point in °C	Boiling point in °C
Alcohol	-98	65
Aluminium	660	2470
Copper	1077	2567
Glycerol	18	290
Lead	328	1737
Iron	1537	2737

Radiation weighting factors

Type of radiation	Radiation weighting factor			
alpha	20			
beta	1			
fast neutrons	10			
gamma	1			
slow neutrons	3			
X-rays	1			

SECTION 1 Attempt ALL questions

1. Two circuits are set up as shown.




Both circuits are used to determine the resistance of resistor R. Which row in the table identifies meter X, meter Y and meter Z?

	meter X	meter Y	meter Z	
А	ohmmeter	voltmeter	ammeter	
В	ohmmeter	ammeter	voltmeter	
С	voltmeter	ammeter	ohmmeter	
D	ammeter	voltmeter	ohmmeter	
Е	voltmeter	ohmmeter	ammeter	

- 2. Which of the following statements is/are correct?
 - I The voltage of a battery is the number of joules of energy it gives to each coulomb of charge.
 - II A battery only has a voltage when it is connected in a complete circuit.
 - III Electrons are free to move within an insulator.
 - A I only
 - B II only
 - C III only
 - D II and III only
 - E I, II and III

3. A circuit is set up as shown.



The resistance between X and Y is


- A 1·3Ω
- B 4·5Ω
- **C** 6.0 Ω
- D 8.0Ω
- E 12 Ω.
- 4. The rating plate on an electrical appliance is shown.



The resistance of this appliance is

- Α 0.017 Ω
- B 0·25 Ω
- C 4·0 Ω
- D 18·4Ω
- E 57·5 Ω.

5. A syringe containing air is sealed at one end as shown.



The piston is pushed in slowly.

There is no change in temperature of the air inside the syringe.

Which of the following statements describes and explains the change in pressure of the air in the syringe?

- A The pressure increases because the air particles have more kinetic energy.
- B The pressure increases because the air particles hit the sides of the syringe more frequently.
- C The pressure increases because the air particles hit the sides of the syringe less frequently.
- D The pressure decreases because the air particles hit the sides of the syringe with less force.
- E The pressure decreases because the air particles have less kinetic energy.
- 6. The pressure of a fixed mass of gas is 150 kPa at a temperature of 27 °C.

The temperature of the gas is now increased to 47 °C.


The volume of the gas remains constant.

The pressure of the gas is now

- A 86 kPa
- B 141 kPa
- C 150 kPa
- D 160 kPa
- E 261 kPa.

[Turn over

7. The diagram represents a water wave.



The wavelength of the water wave is

- A 2 mm
- B 3 mm
- C 4 mm
- D 6 mm
- E 18 mm.
- 8. A student makes the following statements about different types of electromagnetic waves.
 - I Light waves are transverse waves.
 - II Radio waves travel at 340 m s^{-1} through air.
 - III Ultraviolet waves have a longer wavelength than infrared waves.

Which of these statements is/are correct?

- A I only
- B I and II only
- C I and III only
- D II and III only
- E I, II and III
- **9.** Alpha radiation ionises an atom.

Which statement describes what happens to the atom?

- A The atom splits in half.
- B The atom releases a neutron.
- C The atom becomes positively charged.
- D The atom gives out gamma radiation.
- E The atom releases heat.

10. A sample of tissue is irradiated using a radioactive source.

A student makes the following statements.

The equivalent dose received by the tissue is

- I reduced by shielding the tissue with a lead screen
- II increased as the distance from the source to the tissue is increased
- III increased by increasing the time of exposure of the tissue to the radiation.

Which of the statements is/are correct?

- A I only
- B II only
- C I and II only
- D II and III only
- E I and III only
- 11. A sample of tissue receives an absorbed dose of 16μ Gy from alpha particles. The radiation weighting factor for alpha particles is 20.

The equivalent dose received by the sample is

- A 0.80 μSv
- B 1·25 μSv
- C 4 μSv
- D 36 μSv
- E 320 μSv.
- 12. For a particular radioactive source, 240 atoms decay in 1 minute. The activity of this source is
 - A 4 Bq
 - B 180 Bq
 - C 240 Bq
 - D 300 Bq
 - E 14 400 Bq.

[Turn over

13. The letters **X**, **Y** and **Z** represent missing words from the following passage.

During a nuclear X reaction two nuclei of smaller mass number combine to produce a nucleus of larger mass number. During a nuclear Y reaction a nucleus of larger mass number splits into two nuclei of smaller mass number. Both of these reactions are important because these processes can release Z.

	X	Y	Z	
А	fusion	fission	electrons	
В	fission	fusion	energy	
С	fusion	fission	protons	
D	fission	fusion	protons	
E	fusion	fission	energy	

Which row in the table shows the missing words?

- 14. Which of the following quantities is fully described by its magnitude?
 - A Force
 - B Displacement
 - C Energy
 - D Velocity
 - E Acceleration


15. The table shows the velocities of three objects X, Y and Z over a period of 3 seconds. Each object is moving in a straight line.

Time (s)	0	1	2	3
Velocity of X (m s ⁻¹)	2	4	6	8
Velocity of Y (m s ⁻¹)	0	1	2	3
Velocity of Z (m s ⁻¹)	0	2	5	9

Which of the following statements is/are correct?

- I X moves with constant velocity.
- II Y moves with constant acceleration.
- III Z moves with constant acceleration.
- A I only
- B II only
- C I and II only
- D I and III only
- E II and III only
- 16. A car of mass 1200 kg is travelling along a straight level road at a constant speed of $20 \,\mathrm{m\,s^{-1}}$.

The driving force on the car is 2500 N. The frictional force on the car is 2500 N.



The work done moving the car between point X and point Y is

A 0 J

- B 11 800 J
- C 125 000 J
- D 240 000 J
- E 250 000 J.

17. A person sits on a chair which rests on the Earth. The person exerts a downward force on the chair.



Which of the following is the reaction to this force?

- A The force of the chair on the person
- B The force of the person on the chair
- C The force of the Earth on the person
- D The force of the chair on the Earth
- E The force of the person on the Earth
- **18.** A package falls vertically from a helicopter. After some time the package reaches its terminal velocity.


A group of students make the following statements about the package when it reaches its terminal velocity.

- I The weight of the package is less than the air resistance acting on the package.
- II The forces acting on the package are balanced.
- III The package is accelerating towards the ground at $9.8 \,\mathrm{m \, s^{-2}}$.

Which of these statements is/are correct?

- A I only
- B II only
- C III only
- D I and III only
- E II and III only

- 19. The distance from the Sun to Proxima Centauri is 4.3 light years. This distance is equivalent to
 - A $1.4 \times 10^8 \,\mathrm{m}$
 - B $1.6 \times 10^{14} \,\mathrm{m}$
 - C $6.8 \times 10^{14} \,\mathrm{m}$
 - D $9.5 \times 10^{15} \,\mathrm{m}$
 - E $4 \cdot 1 \times 10^{16}$ m.
- **20.** Light from a star is split into a line spectrum of different colours. The line spectrum from the star is shown, along with the line spectra of the elements calcium, helium, hydrogen and sodium.



The elements present in this star are

- A sodium and calcium
- B calcium and helium
- C hydrogen and sodium
- D helium and hydrogen
- E calcium, sodium and hydrogen.

[END OF SECTION 1. NOW ATTEMPT THE QUESTIONS IN SECTION 2 OF YOUR QUESTION AND ANSWER BOOKLET]

ACKNOWLEDGEMENTS

Question 17-Rob Byron/shutterstock.com

_	FOR OFFICIAL USE						-
N5	National Qualification 2015	15				Mark	
X757/75/01		9	Sect	ion 1-		swe	hysics r Grid tion 2
TUESDAY, 5 MAY 9:00 AM – 11:00 AM						577	
Fill in these boxes and re	ad what is printed		Town				
			100011				
Forename(s)	Surnam	ie			Nu	imber (of seat
Date of birth Day Month	Year	Scottish car	ndidate	e number			
Total marks — 110 SECTION 1 — 20 marks Attempt ALL questions. Instructions for the comple SECTION 2 — 90 marks Attempt ALL questions.	etion of Section 1 a	re given on	Page t	wo.			
Reference may be made to to the Relationship Sheet 2 Care should be taken to g to calculations. Write your answers clearly and rough work is provide identify the question num	(757/75/11. ive an appropriate in the spaces prov d at the end of th	number of rided in this is booklet.	signifi bookl If you	cant figu et. Addi use this	ires in tl tional sp space y	ne fina bace fo vou mu	Il answers or answers ost clearly
booklet. You should score Use blue or black ink. Before leaving the examin Invigilator; if you do not, y	through your rough ation room you mus	work wher	oookle	ave writt t to the			
					/	/	©



The questions for Section 1 are contained in the question paper X757/75/02. Read these and record your answers on the answer grid on *Page three* opposite. Use **blue** or **black** ink. Do NOT use gel pens or pencil.


- 1. The answer to each question is **either** A, B, C, D or E. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is only one correct answer to each question.
- 3. Any rough work must be written in the additional space for answers and rough work at the end of this booklet.

Sample Question

The energy unit measured by the electricity meter in your home is the:


- A ampere
- B kilowatt-hour
- C watt
- D coulomb
- E volt.

The correct answer is B-kilowatt-hour. The answer B bubble has been clearly filled in (see below).



Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.



If you then decide to change back to an answer you have already scored out, put a tick (\checkmark) to the **right** of the answer you want, as shown below:







	Α	В	С	D	Е
1	0	0	0	0	0
2	0	0	0	0	0
3	0	0	0	0	0
4	0	0	0	0	0
5	0	0	0	0	0
6	0	0	0	0	0
7	0	0	0	0	0
8	0	0	0	0	0
9	0	0	0	0	0
10	0	0	0	0	0
11	0	0	0	0	0
12	0	0	0	0	0
13	0	0	0	0	0
14	0	0	0	0	0
15	0	0	0	0	0
16	0	0	0	0	0
17	0	0	0	0	0
18	0	0	0	0	0
19	0	0	0	0	0
20	0	0	0	0	0



[BLANK PAGE]

Γ

L

DO NOT WRITE ON THIS PAGE



Page four

MARKS DO NOT WRITE IN THIS MARGIN

[Turn over for Question 1 on Page six

DO NOT WRITE ON THIS PAGE




Page five

3

3

1. A student sets up the following circuit using a battery, two lamps, a switch and a resistor.




(a) Draw a circuit diagram for this circuit using the correct symbols for the components.

(b) Each lamp is rated 2.5 V, 0.50 A.

Calculate the resistance of one of the lamps when it is operating at the correct voltage.

Space for working and answer


* X 7 5 7 7 5 0 1 0 6 *



[Turn over




Page seven





Page eight





Page nine

3. A technician uses pulses of ultrasound (high frequency sound) to detect imperfections in a sample of steel.

The pulses of ultrasound are transmitted into the steel.


The speed of ultrasound in steel is 5200 m s^{-1} .

Where there are no imperfections, the pulses of ultrasound travel through the steel and are reflected by the back wall of the steel.


Where there are imperfections in the steel, the pulses of ultrasound are reflected by these imperfections.

The reflected pulses return through the sample and are detected by the ultrasound receiver.

The technician transmits pulses of ultrasound into the steel at positions X, Y and Z as shown.



The times between the pulses being transmitted and received for positions X and Y are shown in the graph.



Page ten

3.	(coi	ntinue	ed)	MARKS DO NOT WRITE IN THIS
	(a)	(i)	State the time taken between the pulse being transmitted and received at position X.	MARGIN
		(ii)	Calculate the thickness of the steel sample at position X. Space for working and answer	4
	(b)		he graph on the previous page, draw a line to show the reflected e from position Z.	2
	(c)	The	ultrasound pulses used have a period of $4.0\mu s$.	
		(i)	Show that the frequency of the ultrasound pulses is $2 \cdot 5 \times 10^5$ Hz. Space for working and answer	2
		(ii)	Calculate the wavelength of the ultrasound pulses in the steel sample. Space for working and answer	3

Γ




3. (continued)

(d) The technician replaces the steel sample with a brass sample.

The brass sample has the same thickness as the steel sample at position X.

The technician transmits pulses of ultrasound into the brass at position P as shown.



The time between the ultrasound pulse being transmitted and received at position P is greater than the time recorded at position X in the steel sample.

State whether the speed of ultrasound in brass is less than, equal to or greater than the speed of ultrasound in steel.

You must justify your answer.

2

MARKS DO NOT WRITE IN THIS MARGIN



Page twelve

MARKS DO NOT WRITE IN THIS MARGIN

- A science technician removes two metal blocks from an oven. 4. Immediately after the blocks are removed from the oven the technician measures the temperature of each block, using an infrared thermometer. The temperature of each block is 230 °C.

After several minutes the temperature of each block is measured again. One block is now at a temperature of 123°C and the other block is at a temperature of 187 °C.

Using your knowledge of physics, comment on possible explanations for this difference in temperature.


3

[Turn over



Page thirteen

 Diamonds are popular and sought after gemstones. Light is refracted as it enters and leaves a diamond. The diagram shows a ray of light entering a diamond.



- (a) On the diagram, label the angle of incidence i and the angle of refraction r.
- (b) State what happens to the speed of the light as it enters the diamond.
- (c) The optical density of a gemstone is a measure of its ability to refract light.

Gemstones of higher optical density cause more refraction.

A ray of light is directed into a gemstone at an angle of incidence of 45° .

The angle of refraction is then measured.

This is repeated for different gemstones.

Gemstone	Angle of refraction		
А	24·3°		
В	17·0°		
С	27·3°		
D	19·0°		
E	25∙5°		

Diamond is known to have the highest optical density. Identify which gemstone is most likely to be diamond.



1

MARKS DO NOT

1

1


THIS MARGIN

5. (continued)

(d) Diamond is one of the hardest known substances.

Synthetic diamonds are attached to the cutting edges of drill bits for use in the oil industry.

These drill bits are able to cut into rock.



The area of a single cutter in contact with the rock is $1{\cdot}1\times10^{-5}\,m^2.$

When drilling, this cutter is designed to exert a maximum force of $61\,\mathrm{kN}$ on the rock.

Calculate the maximum pressure that the cutter can exert on the rock. Space for working and answer

3

MARKS DO NOT WRITE IN THIS MARGIN



[Turn over

Page fifteen

- 6. A paper mill uses a radioactive source in a system to monitor the thickness MARKS DO NOT of paper. THIS radioactive source rollers paper -700 Geiger-Müller 00 tube counter Radiation passing through the paper is detected by the Geiger-Müller tube. The count rate is displayed on the counter as shown. The radioactive source has a half-life that allows the system to run continuously. (a) State what happens to the count rate if the thickness of the paper decreases. 1 (b) The following radioactive sources are available. **Radioactive Source** Half-life Radiation emitted W 600 years alpha Х 50 years beta Υ 4 hours beta
 - (i) State which radioactive source should be used.

You must explain your answer.

350 years


Ζ

3

gamma



Page sixteen




Page seventeen

7. A ship of mass $5 \cdot 0 \times 10^6$ kg leaves a port. Its engine produces a forward force of $8 \cdot 0 \times 10^3$ N. A tugboat pushes against one side of the ship as shown. The tugboat applies a pushing force of $6 \cdot 0 \times 10^3$ N.

MARKS DO NOT WRITE IN THIS MARGIN

2

2




(a) (i) By scale drawing, or otherwise, determine the size of the resultant force acting on the ship.
 Space for working and answer

(ii) Determine the direction of the resultant force relative to the $8\cdot 0 \times 10^3$ N force.


Space for working and answer



Page eighteen





Page nineteen



Page twenty

8. (continued)





Calculate the acceleration of this trolley between X and Y. *Space for working and answer*


3

MARKS DO NOT WRITE IN THIS MARGIN

[Turn over



Page twenty-one



Page twenty-two

MARKS DO NOT WRITE IN THIS MARGIN

10. Space exploration involves placing astronauts in difficult environments. Despite this, many people believe the benefits of space exploration outweigh the risks.



Using your knowledge of physics, comment on the benefits and/or risks of space exploration.

3



Page twenty-three

[Turn over

[BLANK PAGE]

Г

L

DO NOT WRITE ON THIS PAGE




Page twenty-four

11. Craters on the Moon are caused by meteors striking its surface.



A student investigates how a crater is formed by dropping a marble into a tray of sand.



- (a) The marble has a mass of 0.040 kg.
 - (i) Calculate the loss in potential energy of the marble when it is dropped from a height of 0.50 m.

Space for working and answer

(ii) Describe the energy change that takes place as the marble hits the sand.

[Turn over

MARKS DO NOT WRITE IN THIS MARGIN

3

1



Page twenty-five


(continued) 11.

(b) The student drops the marble from different heights and measures the diameter of each crater that is formed.

The table shows the student's results.

height (m)	diameter (m)				
0.05	0.030				
0.10	0.044				
0.15	0.053				
0.35	0.074				
0.40	0.076				
0.45	0.076				

(i) Using the graph paper below, draw a graph of these results. (Additional graph paper, if required, can be found on Page twenty-eight)






Page twenty-six

MARKS DO NOT WRITE IN THIS MARGIN

3




Page twenty-seven

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORKING

MARKS DO NOT WRITE IN THIS MARGIN

Additional graph paper for Q11 (b) (i)





Page twenty-eight

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORKING

MARKS DO NOT WRITE IN THIS MARGIN



Page twenty-nine

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORKING

MARKS DO NOT WRITE IN THIS MARGIN



Page thirty

[BLANK PAGE]

L

DO NOT WRITE ON THIS PAGE



Page thirty-one

ACKNOWLEDGEMENTS

Question 10 – MarcelClemens/shutterstock.com Question 11 – Procy/shutterstock.com



Page thirty-two



X757/75/11

Physics Relationships Sheet

TUESDAY, 5 MAY 9:00 AM - 11:00 AM





$$E_p = mgh$$
 $d = vt$

$$E_k = \frac{1}{2}mv^2 \qquad \qquad v = f\lambda$$

$$Q = It T = \frac{1}{f}$$

$$V = IR$$

$$A = \frac{N}{2}$$

$$R_T = R_1 + R_2 + \dots \qquad \qquad A = -\frac{1}{t}$$

$$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \dots \qquad D = \frac{E}{m}$$

$$V_2 = \left(\frac{R_2}{R_1 + R_2}\right) V_s \qquad \qquad H = Dw_R$$
$$\dot{H} = \frac{H}{H}$$

$$\frac{V_1}{V_2} = \frac{R_1}{R_2} \qquad \qquad t \qquad \qquad s = vt$$

$$P = \frac{E}{t} \qquad \qquad d = \overline{vt}$$

$$P = IV$$

$$P = I^2 R \qquad \qquad a = \frac{v - u}{t}$$

$$P = \frac{V^2}{R} \qquad \qquad W = mg$$
$$F = ma$$

$$E_h = cm \Delta T \qquad \qquad E_w = Fd$$

$$p = \frac{F}{A} \qquad \qquad E_h = ml$$

$$\frac{pV}{T} = \text{constant}$$
$$p_1 V_1 = p_2 V_2$$

$$\frac{p_1}{T_1} = \frac{p_2}{T_2}$$
$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\overline{T_1} - \overline{T_2}$$

Additional Relationships

Circle

circumference = $2\pi r$

area = πr^2

Sphere

area = $4\pi r^2$

volume = $\frac{4}{3}\pi r^3$

Trigonometry

 $\sin \Theta = \frac{\text{opposite}}{\text{hypotenuse}}$

 $\cos \Theta = \frac{\text{adjacent}}{\text{hypotenuse}}$

 $\tan \Theta = \frac{\text{opposite}}{\text{adjacent}}$

 $\sin^2\theta + \cos^2\theta = 1$

	87 Fr 2,8,18,32, 18,8,1 Francium	55 Cs 2,8,18,18, 8,1 Caesium	37 Rb 2,8,18,8,1 Rubidium	19 K 2,8,8,1 Potassium	Na 2,8,1 Sodium	۲,۰ Lithium 11	² . Γ . ω	Hydrogen	Group 1
La	88 Ra 12, 2,8,18,32, 18,8,2 m Radium	2,8, Bi	38 Sr 3,1 2,8,18,8,2 m Strontium	20 Ca 1 2,8,8,2 m Calcium	Mg 2,8,2 Magnesium	Be	² Be	2n (2)	1 Group 2
Lanthanides Actinides	89 Ac 2,8,18,32, 18,9,2 Actinium	57 La 2,8,18,18, 9,2 Lanthanum	39 Y 2,8,18,9,2 Yttrium	21 Sc 2,8,9,2 Scandium	(3)			_	
57 La 2,8,18, 18,9,2 Lanthanum 89 AC 2,8,18,32, 18,9,2 Actinium	104 Rf 2,8,18,32, 32,10,2 Rutherfordium	72 Hf 2,8,18,32, 10,2 Hafnium	40 Zr 2,8,18, 10,2 Zirconium	22 Ti 2,8,10,2 Titanium	(4)			Key	;
58 Ce 2,8,18, 20,8,2 Cerium 90 Th 2,8,18,32, 18,10,2 Thorium	105 Db 2,8,18,32, 32,11,2 Dubnium	ш, ² , ³⁰	41 Nb 2,8,18, 12,1 Niobium	23 V 2,8,11,2 Vanadium	(5)			Ato	
59 Pr 2,8,18,21, 8,2 Praseodymium 91 91 Pa 2,8,18,32, 20,9,2 Protactinium	106 Sg 2,8,18,32, 32,12,2 Seaborgium	74 W 2,8,18,32, 12,2 Tungsten	42 Mo 2,8,18,13, 1 Molybdenum	24 Cr 2,8,13,1 Chromium	(6) T		Name	Atomic number Symbol Flectron arrangement	
60 Nd 2,8,18,22, 8,2 Neodymium 92 U 2,8,18,32, 21,9,2 Uranium	107 Bh 2,8,18,32, 32,13,2 Bohrium	im 32,	43 Tc 2,8,18,13, 2 Technetium	25 Mn 2,8,13,2 Manganese	(7) (8)			Der] (
61 Pm 2,8,18,23, 8,2 Promethium 93 93 93 93 2,8,18,32, 22,9,2 Neptunium	108 Hs 2,8,18,32, 32,14,2 Hassium	m 32,	44 Ru 2,8,18,15, 1 Ruthenium	26 Fe 2,8,14,2 Iron	Element				
62 Sm 2,8,18,24, 8,2 Samarium 94 94 94 2,8,18,32, 24,8,2 Plutonium	109 Mt 2,8,18,32, 32,15,2 Meitnerium	77 Ir 2,8,18,32, 15,2 Iridium	45 Rh 2,8,18,16, 1 Rhodium	27 Co 2,8,15,2 Cobalt	(6) S				
63 Eu 2,8,18,25, 8,2 Europium 95 Am 2,8,18,32, 2,8,18,32, Americium	110 Ds 2,8,18,32, 32,17,1 Darmstadtium	2,8 Pl	46 Pd 2,8,18, 18,0 Palladium	28 Ni 2,8,16,2 Nickel	(10)				
64 Gd 2,8,18,25, 9,2 Gadolinium 96 Cm 2,8,18,32, 2,25,18,32, Curium	111 Rg 2,8,18,32, 32,18,1 Roentgenium	79 Au 2,8,18, 32,18,1 Gold	47 Ag 2,8,18, 18,1 silver	29 Cu 2,8,18,1 Copper	(11)				
65 Tb 2,8,18,27, 8,2 Terbium 97 97 97 8k 2,8,18,32, 27,8,2 Berkelium	112 Cn 2,8,18,32, 32,18,2 Copernicium	80 Hg 2,8,18, 32,18,2 Mercury	48 Cd 2,8,18, 18,2 Cadmium	30 Zn 2,8,18,2 Zinc	(12)				
66 Dy 2,8,18,28, 8,2 Dysprosium 98 Cf 2,8,18,32, 28,8,2 Californium		81 T 2,8,18, 32,18,3 Thallium	49 In 2,8,18, 18,3 Indium	31 Ga 2,8,18,3 Gallium	Al 2,8,3 Aluminium	Boron 13	ა თ თ	(13)	Group 3
67 Ho 2,8,18,29, 8,2 Holmium 99 Es 2,8,18,32, 29,8,2 Einsteinium		82 Pb 2,8,18, 3 32,18,4 n Lead	50 Sn 18,4 Tin	32 Ge 3 2,8,18,4 1 Germanium	Si 2,8,4 Im Silicon	Carbon 14	°, ∪ ∘	(14)	3 Group 4
68 Er 2,8,18,30, 8,2 Erbium 100 Fm 2,8,18,32, 30,8,2 Fermium		83 Bi 2,8,18, 4 32,18,5 Bismuth	51 Sb , 2,8,18, 18,5 Antimony	33 As 4 2,8,18,5 Jm Arsenic	P 2,8,5 Phosphorus	Nitrogen	² Z V	(15)	4 Group 5
69 Tm 2,8,18,31, 8,2 Thulium 101 101 2,8,18,32, 31,8,32, Mendelevium		84 Po 2,8,18, 5 32,18,6 Polonium	52 Te , 2,8,18, 18,6 ny Tellurium	34 Se 5 2,8,18,6 : Selenium	S 2,8,6 Us Sulfur	n Oxygen	² 0 8	(16)	5 Group 6
70 Yb 2,8,18,32, 8,2 Ytterbium 102 102 2,8,18,32, 32,8,1 Nobelium		85 At 2,8,18, 5 32,18,7 Astatine	53 	2,8 Br	Cl 2,8,7 Chlorine	1 Fluorine 17	ہ <mark>ہ</mark> د	(17)	6 Group 7
71 Lu 2,8,18,32, 9,2 Lutetium 103 Lr 2,8,18,32, 32,9,2 Lawrencium		86 Rn 2,8,18, 32,18,8 Radon	54 Xe 2,8,18, 18,8 Xenon	36 Kr 7 2,8,18,8 Krypton	Ar 2,8,8 Argon	Neon 18	2 Ne	2 He Helium	Gro

Page four

Electron Arrangements of Elements